

The Role of Endoscopy in Diagnosing and Managing Gastroesophageal Reflux Disease (GERD)

Dr. Amit Sachdeva and Nasim Ahmed

¹Assistant Professor, Department of Community Medicine, Indira Gandhi Medical College, Shimla, Himachal Pradesh, India. Email: dramitsachdeva2410@gmail.com

²Independent Research, Guwahati, Assam India. Email: nasim@iarcon.org

Received: Jul. 11, 2025; Revised: Aug. 03, 2025; Accepted: Aug. 22, 2025; Published: Sep. 30, 2025

Under a Creative Commons license Doi: 10.47310/medlet.2025.v02i02.009

Abstract: Gastroesophageal reflux disease (GERD) is a prevalent chronic condition characterized by the reflux of gastric contents into the esophagus, leading to symptoms like heartburn and complications such as esophagitis, Barrett's esophagus, and esophageal adenocarcinoma. Endoscopy, specifically upper gastrointestinal endoscopy, is critical in diagnosing and managing GERD, especially in patients with alarm symptoms, refractory cases, or suspected complications. It allows for direct visualization, biopsy, and therapeutic interventions, including endoscopic fundoplication, radiofrequency ablation, and mucosal resection. Recent advancements, such as narrow band imaging, confocal laser endomicroscopy, and AI-driven diagnostics, have enhanced the detection of early disease and improved patient outcomes. As endoscopic techniques evolve, they offer minimally invasive, personalized treatment options, shaping the future of GERD management.

Keywords: GERD, endoscopy, Barrett's esophagus, esophagitis, radiofrequency ablation, endoscopic fundoplication.

INTRODUCTION

Gastroesophageal reflux disease (GERD) is a common chronic condition that affects millions of people globally. It is characterized by the reflux of stomach contents, particularly acidic gastric juice, into the esophagus, leading to symptoms such as heartburn, regurgitation, and dysphagia. Over time, GERD can result in complications such as esophagitis, Barrett's esophagus, and esophageal adenocarcinoma. Endoscopy, particularly upper gastrointestinal endoscopy (esophagogastroduodenoscopy or EGD), plays a pivotal role in diagnosing and managing GERD.[1-5]

Endoscopy provides a direct view of the esophagus, stomach, and duodenum, allowing clinicians to assess the mucosal surface, detect complications of GERD, and perform therapeutic interventions when necessary. While the diagnosis of GERD is often clinical, especially in uncomplicated cases, endoscopy is crucial in confirming the diagnosis in patients with alarm symptoms, evaluating the severity of esophagitis, detecting complications, and guiding management strategies.[6-8]

This article provides a comprehensive overview of the role of endoscopy in diagnosing and managing GERD, discusses current trends in endoscopic techniques, and explores emerging technologies that are reshaping the field of GERD management.

Overview of GERD [1-5]

1. Pathophysiology of GERD

GERD occurs when the lower esophageal sphincter (LES) fails to maintain an adequate barrier between the stomach and the esophagus. The LES is a ring of muscle located at the junction of the esophagus and stomach, and its relaxation or incompetence allows gastric contents to flow back into the esophagus. The severity of GERD is influenced by factors such as the frequency of reflux events, the volume of refluxed material, and the ability of the esophagus to clear the acidic contents.

Other contributing factors to GERD include impaired esophageal motility, delayed gastric emptying, and increased intra-abdominal pressure, which may occur due to obesity or pregnancy. Lifestyle factors such as

smoking, alcohol consumption, and certain dietary habits (e.g., high-fat meals, chocolate, caffeine) can also exacerbate GERD symptoms.

2. Clinical Presentation and Diagnosis

The typical symptoms of GERD include heartburn (a burning sensation in the chest) and acid regurgitation (the backflow of stomach acid into the throat or mouth). Other symptoms may include chest pain, dysphagia (difficulty swallowing), chronic cough, and hoarseness. GERD can also manifest as extra-esophageal symptoms such as asthma, laryngitis, and dental erosions.

In most patients, GERD is diagnosed based on clinical symptoms and response to proton pump inhibitors (PPIs). However, endoscopy is indicated in patients with alarm features such as dysphagia, odynophagia (painful swallowing), weight loss, anemia, or those with suspected complications like Barrett's esophagus or esophageal adenocarcinoma.

The Role of Endoscopy in Diagnosing GERD [6-8]

1. Indications for Endoscopy in GERD

Endoscopy is not required in all patients with GERD, particularly in those with uncomplicated symptoms that respond well to acid-suppressive therapy. However, endoscopy is essential in the following scenarios:

- **Alarm Symptoms:** Patients with dysphagia, odynophagia, gastrointestinal bleeding, or unintentional weight loss should undergo endoscopy to rule out complications such as esophageal strictures, malignancies, or severe esophagitis.
- **Refractory GERD:** In patients whose symptoms do not improve with standard PPI therapy, endoscopy is used to exclude other causes of symptoms, such as eosinophilic esophagitis, peptic ulcers, or esophageal motility disorders.
- **Assessing Complications:** Endoscopy is crucial for diagnosing complications of GERD, including erosive esophagitis, Barrett's esophagus, esophageal strictures, and esophageal adenocarcinoma.
- **Screening for Barrett's Esophagus:** Patients with long-standing GERD (more than 5 years) and risk factors for Barrett's esophagus (such as obesity, male gender, smoking, or a family history of esophageal cancer) should undergo endoscopy for screening and surveillance.

2. Endoscopic Findings in GERD

The most common findings in GERD during endoscopy include:

- **Erosive Esophagitis:** Endoscopy can reveal varying degrees of esophageal inflammation, ranging from mild erythema and friability to severe ulcerations and erosions. The Los Angeles (LA) classification system is commonly used to grade the severity of esophagitis, with grades ranging from A (mild) to D (severe).
- **Barrett's Esophagus:** This premalignant condition is characterized by the replacement of normal squamous epithelium in the esophagus with metaplastic columnar epithelium. Endoscopy allows for the visualization of this change, and biopsies are typically obtained to confirm the diagnosis and assess for dysplasia.
- **Esophageal Strictures:** These are areas of narrowing in the esophagus due to fibrosis and scarring from chronic inflammation. Endoscopy is used to visualize strictures and perform dilation if necessary.
- **Esophageal Adenocarcinoma:** Although rare, GERD is a known risk factor for esophageal adenocarcinoma. Endoscopy allows for early detection of malignancies and biopsy of suspicious lesions.

3. Endoscopic Biopsy and Histopathology

In addition to visualizing the esophageal mucosa, endoscopy allows for tissue sampling (biopsy), which is essential for confirming the presence of complications like Barrett's esophagus and ruling out malignancies or other causes of esophageal inflammation. Biopsy specimens can reveal histopathological features such as:

- **Inflammatory Changes:** In GERD, biopsies often show basal cell hyperplasia, elongation of the papillae, and inflammatory cell infiltration.
- **Dysplasia in Barrett's Esophagus:** Biopsies from areas of Barrett's esophagus are analyzed for the presence of dysplasia, which is a precursor to esophageal adenocarcinoma. Surveillance endoscopy with biopsies is recommended to detect dysplasia early and guide treatment decisions.

Endoscopy in the Management of GERD [9-12]

1. Endoscopic Therapy for GERD

While most GERD patients are managed with lifestyle modifications and pharmacotherapy, a subset of patients with refractory symptoms or those seeking alternatives to long-term PPI use may benefit from endoscopic therapies. These procedures aim to enhance the function of the LES and reduce acid reflux. Common endoscopic treatments include:

Endoscopic Fundoplication (TIF)

Transoral incisionless fundoplication (TIF) is an endoscopic procedure that replicates the effects of surgical fundoplication without the need for incisions. During TIF, the esophagus is accessed through the mouth, and a specialized device is used to create a partial wrap of the gastric fundus around the lower esophagus, enhancing the pressure of the LES and preventing reflux.

Studies have shown that TIF can improve GERD symptoms, reduce PPI dependence, and improve quality of life. However, its long-term efficacy is still being evaluated, and it may not be suitable for patients with large hiatal hernias.

Radiofrequency Ablation (Stretta)

The Stretta procedure involves the delivery of radiofrequency energy to the muscles of the LES and gastric cardia. This energy induces controlled scarring, leading to the thickening of the LES and reduction of reflux episodes.

While some studies have reported improvements in GERD symptoms following the Stretta procedure, its use remains controversial, with some clinical guidelines recommending caution due to inconsistent long-term results.

2. Endoscopic Management of Complications

Endoscopy is crucial not only for diagnosing GERD-related complications but also for managing them:

Dilation of Esophageal Strictures

Esophageal strictures can cause progressive dysphagia and significantly impair quality of life. Endoscopic dilation, using either balloon or bougie dilators, is an effective and safe method for relieving esophageal strictures. In some cases, multiple sessions may be required to achieve optimal results.

Endoscopic Mucosal Resection (EMR)

For patients with Barrett's esophagus and early esophageal adenocarcinoma, endoscopic mucosal resection (EMR) is a minimally invasive alternative to surgery. During EMR, the superficial layers of the esophagus, containing the abnormal or cancerous tissue, are removed endoscopically. This procedure allows for the removal of dysplastic or early neoplastic tissue without the need for more extensive surgery.

Radiofrequency Ablation (RFA) for Barrett's Esophagus

In patients with Barrett's esophagus and low-grade or high-grade dysplasia, radiofrequency ablation (RFA) is used to destroy the dysplastic cells. During RFA, a catheter-mounted balloon is passed through the endoscope, and radiofrequency energy is delivered to the affected areas of the esophagus, ablating the abnormal tissue.

RFA has been shown to be highly effective in eradicating dysplasia and reducing the risk of progression to esophageal adenocarcinoma. In combination with surveillance endoscopy, RFA is an important tool in the management of Barrett's esophagus.

Current Trends and Emerging Endoscopic Technologies in GERD [12-15]

1. Advanced Imaging Techniques

Endoscopic technologies have evolved significantly, improving the ability to detect subtle lesions and early-stage disease. Advanced imaging techniques include:

Narrow Band Imaging (NBI)

Narrow band imaging (NBI) enhances the visualization of the mucosal surface and vasculature by using specific wavelengths of light. This technique improves the detection of early mucosal changes in Barrett's esophagus and other GERD-related conditions, facilitating early diagnosis and intervention.

Confocal Laser Endomicroscopy (CLE)

Confocal laser endomicroscopy (CLE) is an advanced imaging technique that allows for real-time, in vivo microscopic evaluation of the esophageal mucosa. CLE can differentiate between normal and dysplastic tissue at a cellular level, enhancing the accuracy of biopsy targeting in Barrett's esophagus.

2. Artificial Intelligence (AI) in Endoscopy

Artificial intelligence (AI) is revolutionizing the field of endoscopy, with the development of AI-powered systems that can assist in the detection and diagnosis of GERD-related lesions. AI algorithms, trained on large datasets of endoscopic images, can detect abnormalities such as esophagitis, Barrett's esophagus, and early-stage cancer with high accuracy.

AI has the potential to reduce interobserver variability, improve diagnostic accuracy, and enhance the efficiency of endoscopic procedures. As AI technology continues to advance, it is likely to play an increasingly important role in GERD management.

The Future of Endoscopy in GERD [11-16]

As the understanding of GERD continues to evolve, so too will the role of endoscopy in its diagnosis and management. Emerging technologies and therapeutic innovations are reshaping the landscape of GERD treatment, offering new opportunities to enhance patient care.

Endoscopic Anti-Reflux Devices

Several novel endoscopic devices are under investigation for the treatment of GERD. These devices aim to restore the function of the LES and prevent reflux without the need for surgery. Examples include the magnetic sphincter augmentation device (LINX), which uses a ring of magnetic beads to reinforce the LES, and endoscopic suturing devices that tighten the LES.

Personalized Endoscopic Approaches

The future of GERD management may involve personalized approaches, in which endoscopic findings and patient characteristics are used to tailor treatment strategies. For example, patients with mild GERD and no complications may be managed conservatively with lifestyle modifications and PPIs, while those with refractory GERD or high-risk Barrett's esophagus may benefit from endoscopic interventions.

CONCLUSION

Endoscopy plays an essential role in both the diagnosis and management of gastroesophageal reflux disease. It is indispensable for identifying complications such as esophagitis, Barrett's esophagus, and esophageal cancer, and for guiding treatment strategies in patients with refractory symptoms or those at risk for progression. In addition, endoscopic therapies, such as TIF, radiofrequency ablation, and endoscopic resection, offer minimally invasive alternatives to surgery, improving outcomes for patients with GERD and its complications.

As technology continues to advance, the integration of AI, advanced imaging techniques, and novel therapeutic devices will further enhance the role of endoscopy in GERD management. With these innovations, endoscopy is set to remain at the forefront of GERD care, providing patients with more effective, personalized, and less invasive treatment options.

REFERENCES

1. Clarrett et al; "Gastroesophageal Reflux Disease (GERD)" Missouri Medicine 115.3 (2018) Pp 214-218
2. Mayo Clinic; "Gastroesophageal Reflux Disease (GERD): Symptoms and Causes" Mayo Clinic (n.d.) Pp n/a, available from <https://www.mayoclinic.org/diseases-conditions/gerd/symptoms-causes/syc-20361940>
3. MedlinePlus; "Gastroesophageal Reflux Disease" MedlinePlus (n.d.) Pp n/a, available from <https://medlineplus.gov/ency/article/000265.htm>
4. WebMD; "Reflux Disease (GERD)" WebMD (n.d.) Pp n/a, available from <https://www.webmd.com/heartburn-gerd/reflux-disease-gerd-1>
5. Physio-pedia; "Gastroesophageal Reflux Disease" Physio-pedia (n.d.) Pp n/a, available from https://www.physio-pedia.com/Gastroesophageal_Reflux_Disease
6. Kuribayashi et al; "The Role of Endoscopy in the Management of Gastroesophageal Reflux Disease" DEN Open 2.1 (2021) Pp e86, doi <https://doi.org/10.1002/deo2.86>
7. Simadibrata et al; "Role of Endoscopy in Gastroesophageal Reflux Disease" Clinical Endoscopy 56.6 (2023) Pp 681-692, doi <https://doi.org/10.5946/ce.2023.182>
8. Cleveland Clinic; "Esophagogastroduodenoscopy (EGD) Test" Cleveland Clinic (n.d.) Pp n/a, available from <https://my.clevelandclinic.org/health/diagnostics/22549-esophagogastroduodenoscopy-egd-test>
9. Lopes et al; "Endoscopic Treatment of Gastroesophageal Reflux: A Narrative Review" Porto Biomedical Journal 8.4 (2023) Pp e226, doi <https://doi.org/10.1097/j.pbj.0000000000000226>
10. Lee et al; "Endoscopic Management of GERD" Digestive Diseases and Sciences 67 (2022) Pp 1455-1468, doi <https://doi.org/10.1007/s10620-021-07380-x>

11. Akcelik et al; "Endoscopic Interventions in the Management of the Gastroesophageal Reflux: A Narrative Review" *Annals of Esophagus* 6 (2023) Pp 23, doi <https://doi.org/10.21037/aoe-22-25>
12. Healthline; "GERD Endoscopy" Healthline (n.d.) Pp n/a, available from <https://www.healthline.com/health/gerd-endoscopy>
13. Calabrese et al; "New Perspectives in Endoscopic Treatment of Gastroesophageal Reflux Disease" *Diagnostics (Basel)* 13.12 (2023) Pp 2057, doi <https://doi.org/10.3390/diagnostics13122057>
14. Loganathan et al; "Endoscopic Advances in the Diagnosis and Management of Gastroesophageal Reflux Disease" *Medicina* 60.7 (2024) Pp 1120, doi <https://doi.org/10.3390/medicina60071120>
15. Mann et al; "Advanced Endoscopic Imaging and Interventions in GERD: An Update and Future Directions" *Frontiers in Medicine (Lausanne)* 8 (2021) Pp 728696, doi <https://doi.org/10.3389/fmed.2021.728696>
16. Simadibrata et al; "Role of Endoscopy in Gastroesophageal Reflux Disease" *Clinical Endoscopy* 56.6 (2023) Pp 681-692, doi <https://doi.org/10.5946/ce.2023.182>